VOL. 443 JOURNAL OF CHROMATOGRAPHY JUNE 29, 1988

CONTENTS

7TH INTERNATIONAL SYMPOSIUM ON HIGH-PERFORMANCE LIQUID CHROMATOGRAPHY OF PROTEINS, PEPTIDES AND POLYNUCLEOTIDES, WASHINGTON, DC, NOVEMBER 2–4, 1987, PART I

Foreword
by C. T. Wehr 1

High-performance immobilized-metal-ion affinity chromatography of peptides and proteins
by J. Porath (Uppsala, Sweden) 3

Preparative chromatography of proteins. Analysis of the multivalent ion-exchange formalism
by A. Velayudhan and Cs. Horváth (New Haven, CT, U.S.A.) 13

Tandem separation schemes for preparative high-performance liquid chromatography of proteins
by A. L. Lee, A. W. Liao and Cs. Horváth (New Haven, CT, U.S.A.) 31

Comparative study of Zorbax Bio Series GF 250 and GF 450 and TSK-Gel 3000 SW and SWXL columns in size-exclusion chromatography of proteins
by B. Anspach, H. U. Gierlich and K. K. Unger (Mainz, F.R.G.) 45

Dextran-coated silica packings for high-performance size-exclusion chromatography of proteins
by X. Santarelli, D. Muller and J. Jozefonvicz (Villetaneuse, France) 55

Peptide characterization with a sulfoethyl aspartamide column
by D. L. Crimmins, J. Gorka, R. S. Thoma and B. D. Schwartz (St. Louis, MO, U.S.A.) 63

Synthesis of a non-porous, polystyrene-based strong anion-exchange packing material and its application to fast high-performance liquid chromatography of proteins
by M. A. Rounds and F. E. Regnier (West Lafayette, IN, U.S.A.) 73

Cation-exchange chromatography of peptide on poly(2-sulfoethyl aspartamide)-silica
by A. J. Alpert (Columbia and Baltimore, MD, U.S.A.) and P. C. Andrews (West Lafayette, IN, U.S.A.) 85

High-performance liquid chromatography of amino acids, peptides and proteins. LXXXVI. The influence of different displacer salts on the retention and bandwidth properties of proteins separated by isocratic anion-exchange chromatography
by M. T. W. Hearn, A. N. Hodder and M. I. Aguilar (Clayton, Australia) 97

Molecular orientation of immunoglobulin G at high concentration on an ion-exchange sorbent
by I. Mazsaroff, S. Cook and F. E. Regnier (West Lafayette, IN, U.S.A.) 119

Ion-exchange chromatography of proteins on a polyethyleneimine-grafted hydrophylic polymer for high-performance liquid chromatography
by N. Kitagawa (Mountain View, CA, U.S.A.) 133

Chromatographic evaluation of large-pore and non-porous polymeric reversed phases
by D. P. Lee (Reno, NV, U.S.A.) 143

Studies on the stabilization of reversed phases for liquid chromatography
by N. Sagliano, Jr., T. R. Floyd and R. A. Hartwick (Piscataway, NJ, U.S.A.) and J. M. DiBussolo and N. T. Miller (Lafayette Hill, PA, U.S.A.) 155

Chromatography of proteins on hydrophobic interaction and ion-exchange chromatographic matrices: mobile phase contributions to selectivity
by M. L. Heinitz (Minneapolis, MN, U.S.A.) and L. Kennedy, W. Kopaciewicz and F. E. Regnier (West Lafayette, IN, U.S.A.) 173

Reduction studies on bacterial recombinant somatomedin C/insulin-like growth factor-I
by H. Meng, B. D. Burleigh and G. M. Kelly (Northbrook, IL, U.S.A.) 183

Surface-mediated retention effects of subtilisin site-specific variants in cation-exchange chromatography
by R. M. Chicz and F. E. Regnier (West Lafayette, IN, U.S.A.) 193

Reversible subunit dissociation of tumor necrosis factor during hydrophobic interaction chromatography
by M. G. Kunitani and R. L. Cunico (Emeryville, CA, U.S.A.) and S. J. Staats (Seattle, WA, U.S.A.) 205

Chromatographic resolution of lysozyme variants
by J. Fausnaugh-Pollitt, G. Thevenon, L. Janis and F. E. Regnier (West Lafayette, IN, U.S.A.) 221

Correlation of calcitonin structure with chromatographic retention in high-performance liquid chromatography
by M. L. Heinitz (Minneapolis, MN, U.S.A.), E. Flanigan and R. C. Orlowski (Kankakee, IL, U.S.A.) and F. E. Regnier (West Lafayette, IN, U.S.A.) 229

Effects of reductive alkylation on peptide retention times
by E. M. Brown, H. J. Dower and R. Greenberg (Philadelphia, PA, U.S.A.) 247

Effect of detergents on the structure of integral membrane proteins of Sendai virus studied with size-exclusion high-performance liquid chromatography and monoclonal antibodies
by S. Welling-Wester and B. Kazemier (Groningen, The Netherlands), C. Örvell (Stockholm, Sweden) and G. W. Welling (Groningen, The Netherlands) 255

Structural characterization of glycoproteins
by F. W. Putnam, and N. Takahashi (Bloomington, IN, U.S.A.) 267

Application of high-performance liquid chromatography in neurophysin disulfide assignment
by S. Burman, E. Breslow, B. T. Chait and T. Chaudhary (New York, NY, U.S.A.) . . 285

Determination of monosaccharides in glycoproteins by reversed-phase high-performance liquid chromatography on 2.1-mm narrowbore columns
by D. J. Gisch and J. D. Pearson (Kalamazoo, MI, U.S.A.) 299

Isolation and comparative peptide mapping of fibrinogen subunits by reversed-phase high-performance liquid chromatography
by R. F. Ebert (Rockville, MD, U.S.A.) and C. H. Schmelzer (Baltimore, MD, U.S.A.) . 309

High-performance liquid chromatography of complex mixtures of cyanogen bromide-produced peptides from different proteins
by P. Iadarola, M. C. Zapponi, M. Stoppini, M. L. Meloni, L. Minchiotti, M. Galliano and G. Ferri (Pavia, Italy) 317

High-performance liquid chromatographic method for the determination of ε-(γ-glutamyl)lysine and mono- and bis-γ-glutamyl derivatives of putrescine and spermidine
by S. Beninati, N. Martinet and J. E. Folk (Bethesda, MD, U.S.A.) 329

Single-step purification of two hyperglycaemic neurohormones from the sinus gland of *Procambarus bouvieri*. Comparative peptide mapping by means of high-performance liquid chromatography
by A. Huberman and M. B. Aguilar (México, México) 337

Rapid peptide mapping by high-performance liquid chromatography
by K. Kalghatgi and Cs. Horváth (New Haven, CT, U.S.A.) 343

Microbore high-performance liquid chromatography–mass spectrometry for the analysis of proteolytic digests by continuous-flow fast-atom bombardment mass spectrometry
by R. M. Caprioli, W. T. Moore, B. DaGue and M. Martin (Houston, TX, U.S.A.) . . 355

Separation of proteins by reversed-phase high-performance liquid chromatography. I. Optimizing the column
by W. G. Burton, K. D. Nugent, T. K. Slattery and B. R. Summers (Foster City, CA, U.S.A.) and L. R. Snyder (Orinda, CA, U.S.A.) 363

Separation of proteins by reversed-phase high-performance liquid chromatography. II. Optimizing sample pretreatment and mobile phase conditions
by K. D. Nugent, W. G. Burton, T. K. Slattery and B. F. Johnson (Foster City, CA, U.S.A.) and L. R. Snyder (Orinda, CA, U.S.A.) 381

Author Index 399

In articles with more than one author, the name of the author to whom correspondence should be addressed is indicated in the article heading by a 6-pointed asterisk (*)